

*Euglandina*, a few specimens of one of the small glassy sections.

*Practicolella*, quite good set.

*Polygyra*, infrequent, but must have quite a few.

*Trichodiscina* with animal, I think 2 species.

*Thysanophora*, *T. plagiptycha* at least recognized, on ferns about a foot above ground.

*Drymaeus*, only two species.

*Cælocentrum*, two species, I think.

I wrote you that I spotted F. & C.'s type locality of *Lysinoe humboldtiana* and found a nice electric line that goes right close by. Spent an hour at La Vesta in a torrential rainstorm and got a number of specimens of a big *Drymaeus*, but did not get out to El Desierto. I hope to be enough recuperated by the higher, drier climate of Mexico City so that I can get in some good work. I am also going to make another try for *Leptarionta guillarmodi* and *Ceres* at Cordoba if I can. I will leave Vera Cruz Sept. 11, which will get me into Philadelphia about the 20th. Hoping to see you in about a month, I remain,

Yours truly,

HORACE B. BAKER.

Care of Mexican Light & Power Co.,  
Necaxa, Estado Puebla,  
Ee. Un. Mexico."

---

SOME NOTES AND A LIST OF SHELLS OF RIO, KENTUCKY

---

BY WILLIAM J. CLENCH

---

(Concluded from p. 12)

The following is a complete list of the species obtained at Rio. Where the year is noted it designates that the species was only taken at that time. Where no yearly citation appears, the species was collected both years. This only applies to the Unionidae and Pleuroceridae, as little or no time was spent in 1925 for land shells.

I am under great obligation to Dr. Walker, Dr. Ortmann and Mr. Calvin Goodrich, who have identified most of the material.

Glenbrook Spring	<i>Tritogonia verrucosa</i> (Raf.)
<i>Lymnaea humilis modicella</i>	<i>Cyclonaias tuberculatus</i> Raf.
(Say)	<i>Pleurobema cordatum</i> (Raf.)
<i>Physa microstoma</i> Hald.	1925
<i>Goniobasis laqueata</i> Say	<i>Pleurobema cordatum coccinea</i>
* * *	(Conr.) 1925
About the Spring	<i>Pleurobema cordatum catillus</i>
<i>Polygyra plicata</i> Say	(Conr.) 1925
" <i>tridentata</i> (Say)	<i>Amblema costata</i> Raf.
" <i>inflecta</i> (Say)	<i>Fusconaia subrotunda</i> (Lea)
" <i>albolabris</i> (Say)	<i>Fusconaia subrotunda kirt-</i>
" <i>zaleta</i> (Binn)	<i>landiana</i> (Lea)
" <i>appressa</i> (Say)	<i>Fusconaia ebena</i> (Lea) 1924
" <i>thyroides</i> (Say)	" <i>flava</i> (Raf.)
" <i>stenotrema</i> (Fer.)	<i>Megalonaias gigantea</i> (Barnes)
<i>Gastrocopta armifera</i> Say	1925
<i>Mesophix laevigata</i> Pfr.	<i>Elliptio crassidens</i> (Lam)
<i>Vitrea indentata</i> Say	" <i>dilatatus</i> (Raf.)
" <i>capsella</i> (Gld.)	<i>Lastena lata</i> (Raf.) 1925
<i>Gastrodonta ligera</i> (Say)	<i>Lasmigona costata</i> (Raf.)
* * *	<i>Alasmidonta marginata</i> (Say)
Green River, at Toll bridge	1925
<i>Campeloma integrum</i> (Say) ?	<i>Strophitus rugosus</i> (Swains)
<i>Pleurocera undulatum</i> Say	1924
<i>Lithasia obovata</i> Say	<i>Ptychobranthus fasciolare</i>
<i>Lampsilis anodontoides</i> (Lea)	(Raf.)
1924	<i>Cyprogenia irrorata</i> (Lea) 1925
<i>Proptera alata</i> (Say) 1924	<i>Obovaria subrotunda</i> (Raf.)
* * *	1924
Green River on the Shoals	<i>Obovaria subrotunda lens</i>
<i>Goniobasis curreyana</i> Lea	(Lea) 1924
<i>Quadrula pustulosa</i> Lea	<i>Actinonaias carinata</i> (Barnes)
" <i>metanevra wardi</i>	<i>Ligumia recta latissima</i> (Raf.)
(Lea)	<i>Eurynia vanuxemensis</i> (Lea)
	1924

Leptodea fragilis (Raf.)	Lampsilis ovata ventricosa
Proptera alata (Say)	(Barnes)
Micromya ortmanni Walker	Truncilla truncata Raf.
Lampsilis siliquoidea (Barnes)	Dysnomia triquetra (Raf.)
1924	1925
Lampsilis ovata (Say) 1925	

## BIBLIOGRAPHY

- 1895—Cooke, A. H. Camb. Nat. Hist. Mollusca.  
 1920—Grier, N. M. NAUTILUS, Vol. XXXIV.  
 1922—Jewell, M. E. Ecology, Vol. III.

---

 A NEW PECTEN FROM THE LOOCHOO ISLANDS
 

---

BY W. H. DALL

---

## PECTEN (CHLAMYS) SMEGMATUS n. sp.

Shell compressed, glistening, whitish above with reddish flecks, darker and more crowded distally; a few opaque white zigzag markings near the umbo; lower valve white with purplish flush near the umbo and reddish margin on the wings; sculpture of the upper valve with nine low subangular whitish rays with one smaller ray in the interspaces with one or two small threads on each side of it; wings nearly smooth, oblique, the left wing much the largest; sculpture of the lower valve of numerous low undulated riblets with one or two small threads in the interspaces; the sculpture grows feeble toward the umbo; byssal notch wide and deep, ctenolium with eleven denticulations; height of shell, 33; width of shell, 30; of hinge-line, 17; diameter, 7 mm.

Loochoo Islands; Coll. Thaanum, No. 9455.

This is nearest some of the forms of *Pecten squamosus* Gmelin, but has a different sculpture with no indication of raised scales. The surface has a greasy sheen. A single specimen was obtained by Messrs. Thaanum and Langford, during their collecting trip of 1925.